POST-TEST

Investigator Perspectives on the Prevalence and Clinical Implications of Inaccurate or Misclassified Lymphoma Diagnoses in Community Oncology Practice

QUESTIONS (PLEASE CIRCLE ANSWER):

1.	The	majority	of	patients	with	del(17p	o) CLL
----	-----	----------	----	----------	------	---------	--------

- a. Present up front with the 17p deletion
- b. Acquire the 17p deletion over the course of their disease
- 2. Which of the following categories reflects the mechanism of action of venetoclax?
 - a. Antibody-drug conjugate
 - b. Anti-PD-1/PD-L1 antibody
 - c. Bcl-2 inhibitor
 - d. PI3K inhibitor
- 3. Treatment with ______ for patients with CLL can result in high rates of MRD negativity, which typically correlates with longer progression-free survival, time to next treatment and longer durations of remission.
 - a. Ibrutinib
 - b. Venetoclax
 - c. Both a and b
 - d. Neither a nor b
- 4. The Phase III ECHELON-1 trial evaluating brentuximab vedotin with doxorubicin, vinblastine and dacarbazine versus bleomycin with doxorubicin, vinblastine and dacarbazine as front-line therapy for patients with previously untreated Stage III or IV HL demonstrated superior _____ for patients who received the brentuximab vedotin-based combination.
 - a. Progression-free survival
 - b. Overall survival
 - c. Both a and b
 - d. Neither a nor b
- 5. Hospitalization for the purpose of monitoring for TLS is required for all patients starting therapy with venetoclax.
 - a. True
 - b. False

- A study published by Bartlett and colleagues evaluating brentuximab vedotin monotherapy in patients with DLBCL reported response rates of 30% to 40% for patients with CD30-undetectable disease.
 - a. True
 - b. False
- 7. Venetoclax is active in patients with
 - a. CLL with 17p deletion
 - b. CLL without 17p deletion
 - c. Both a and b
 - d. Neither a nor b
- 8. A publication by Zinzani and colleagues in *Blood* stated that pembrolizumab monotherapy demonstrated activity in patients with primary mediastinal large B-cell lymphoma.
 - a. True
 - b. False
- 9. Technical issues with core needle biopsies that can lead to inaccurate diagnosis include
 - a. Crush artifact
 - b. Edge effect
 - c. Limited amount of tissue
 - d. All of the above
 - e. Both a and b
 - f. Both a and c
- 10. Which of the following lymphomas exhibits constant expression of the CD30 antigen?
 - a. Anaplastic large cell lymphoma
 - b. DLBCL
 - c. HL
 - d. PTCI