

Cases from the Community: Investigators Discuss Emerging Research and Actual Patients with Renal Cell Carcinoma**THE CORRECT ANSWER IS INDICATED WITH YELLOW HIGHLIGHTING.**

- Alterations in which of the following genes located on chromosome 3 are commonly found in patients with sporadic clear cell renal cell carcinoma (RCC)?**
 - Bcl-2
 - p53
 - VHL (von Hippel-Landau gene)**
- Which of the following combinations were compared to sunitinib as first-line therapy for patients with advanced RCC in the Phase III CLEAR trial?**
 - Lenvatinib with everolimus or pembrolizumab**
 - Tivozanib with everolimus or pembrolizumab
 - Pazopanib with nivolumab or pembrolizumab
- Which of the following drug types best describes the mechanism of action of belzutifan (MK-6482)?**
 - VEGFR tyrosine kinase inhibitor
 - Hypoxia-inducible factor-2 alpha (HIF-2 α) inhibitor**
 - Immune checkpoint inhibitor
- The CABOSUN trial comparing cabozantinib to sunitinib as initial therapy for intermediate- or poor-risk advanced RCC reported which of the following outcomes in terms of progression-free survival?**
 - Improvement with cabozantinib**
 - Improvement with sunitinib
 - Similar progression-free survival in both study arms