POST-TEST

Cases from the Community: Urologic Oncology Investigators Provide Perspectives on the Optimal Management of Urothelial Bladder Cancer (Webinar Video Proceedings)

THE CORRECT ANSWER IS INDICATED WITH YELLOW HIGHLIGHTING.

- 1. Which of the following is targeted by erdafitinib?
 - a. FGFR
 - b. Nectin-4
 - c. Trop-2
 - d. HER3
- 2. The ongoing Phase III POTOMAC trial is evaluating the anti-PD-L1 antibody durvalumab in which of the following settings?
 - a. As monotherapy for patients with high-risk non-muscle-invasive bladder cancer (NMIBC) unresponsive or refractory to BCG therapy
 - In combination with BCG for patients with high-risk, BCG-naïve NMIBC
 - c. As maintenance therapy after frontline chemotherapy for patients with metastatic urothelial bladder carcinoma (mUBC)
 - d. In combination with enfortumab vedotin for patients with mUBC

- 3. Which of the following Grade 3 or higher adverse events was most commonly reported with enfortumab vedotin in the EV-301 trial?
 - a. Diarrhea
 - b. Alopecia
 - c. Febrile neutropenia
 - d. Rash
- 4. What is the mechanism of action of TAR-200?
 - a. Antibody-drug conjugate
 - b. FGFR inhibitor
 - c. Anti-PD-1/PD-L1 antibody
 - d. Intravesicular gemcitabine
- 5. Which of the following any-grade adverse events was most commonly observed in the TROPHY-U-01 trial evaluating sacituzumab govitecan for patients with mUBC?
 - a. Febrile neutropenia
 - b. Diarrhea
 - c. Urinary tract infection
 - d. Dyschromia