

PARP Inhibition in the Management of Prostate Cancer — Where We Are and Where We're Headed

THE CORRECT ANSWER IS INDICATED WITH YELLOW HIGHLIGHTING.

- 1. Patients with which of the following biomarker statuses experience the most durable responses to PARP inhibitors for prostate cancer?**
 - a. BRCA1/2 heterozygous deletion
 - b. BRCA2 homozygous deletion**
 - c. BRCA reversion mutations
 - d. BRCA status does not correlate with duration of response
- 2. Which of the following Grade 3 or higher adverse events was most commonly associated with niraparib in combination with abiraterone acetate and prednisone among patients with metastatic castration-resistant prostate cancer (mCRPC) and specified homologous recombination repair (HRR) gene alterations (HRR biomarker-positive) in the Phase III MAGNITUDE trial?**
 - a. Anemia**
 - b. Cardiac failure
 - c. Cerebrovascular disorder
 - d. Immune-related hepatitis
 - e. Rash
- 3. Results of the Phase III PROpel trial evaluating first-line olaparib in combination with abiraterone versus abiraterone alone included an improvement in radiographic progression-free survival (rPFS) for which patients with mCRPC?**
 - a. Only those with HRR gene mutations
 - b. Only those with no HRR gene mutations
 - c. All patients regardless of HRR gene mutation status**
 - d. No patients regardless of HRR gene mutation status
- 4. The Phase III MAGNITUDE trial evaluating niraparib with abiraterone acetate and prednisone demonstrated a clinically significant benefit for which patients with HRR biomarker-positive status?**
 - a. Those with metastatic hormone-sensitive prostate cancer (mHSPC) receiving first-line therapy
 - b. Those with heavily pretreated mHSPC
 - c. Those with mCRPC receiving first-line therapy**
 - d. Those with heavily pretreated mCRPC
- 5. Results of the Phase III MAGNITUDE trial evaluating first-line niraparib in combination with abiraterone versus abiraterone alone included an improvement in rPFS for which patients with mCRPC?**
 - a. Only those with BRCA1/2 or HRR gene mutations**
 - b. Only those with no BRCA1/2 or HRR gene mutations
 - c. All patients regardless of BRCA or HRR gene mutation status
 - d. No patients regardless of BRCA or HRR gene mutation status