

Meet The Professor: Optimizing the Management of Acute Myeloid Leukemia and Myelodysplastic Syndromes — Part 2 of a 3-Part Series

THE CORRECT ANSWER IS INDICATED WITH YELLOW HIGHLIGHTING.

- In the ASCERTAIN (ASTX727-02) trial for patients with acute myeloid leukemia (AML), the safety profile of oral decitabine/cedazuridine is best characterized by which statement below?**
 - The safety profile was consistent with that of decitabine
 - Most Grade 3 or higher events were related to myelosuppression
 - Most gastrointestinal system adverse events were generally Grade 3 or 4
 - Both a and b**
 - Both b and c
- The Phase III COMMANDS trial evaluating luspatercept versus epoetin alfa for lower-risk transfusion-dependent myelodysplastic syndromes demonstrated which of the following outcomes?**
 - Epoetin alfa was superior to luspatercept in increasing the likelihood of achieving transfusion independence and increasing hemoglobin levels
 - Luspatercept was superior to epoetin alfa in increasing the likelihood of achieving transfusion independence and increasing hemoglobin levels**
 - No difference was observed between luspatercept and epoetin alfa in subsequent transfusion independence or increased hemoglobin levels
- Data from the QuANTUM-First trial led to the recent FDA approval of quizartinib in combination with chemotherapy for patients with AML in which setting?**
 - Newly diagnosed AML with an IDH1 mutation
 - Relapsed AML with an IDH1 mutation
 - Newly diagnosed AML with a FLT3-ITD mutation**
 - Relapsed AML with a FLT3-ITD mutation
- The Phase III placebo-controlled QuANTUM-First trial of quizartinib with chemotherapy for AML demonstrated which of the following outcomes?**
 - A nonsignificant increase in overall survival (OS) with quizartinib/chemotherapy
 - A statistically significant doubling of OS with quizartinib/chemotherapy**
 - No difference in OS between quizartinib/chemotherapy and placebo
- Longer-term follow-up of the placebo-controlled AGILE study of the addition of ivosidenib to azacitidine for newly diagnosed AML demonstrated which outcome?**
 - The OS benefit with ivosidenib/azacitidine was not maintained
 - The OS benefit with ivosidenib/azacitidine was maintained**
 - The proportion of patients converting to transfusion independence was similar to that seen with placebo
 - The proportion of patients converting to transfusion independence was smaller than that seen with placebo