POST-TEST

Striving for Consensus: Current and Future Management of Metastatic Non-Small Cell Lung Cancer in Patients Who Experience Disease Progression on Immune Checkpoint Inhibitor Therapy

THE CORRECT ANSWER IS INDICATED WITH YELLOW HIGHLIGHTING.

- 1. Which of the following is targeted by antibody-drug conjugate datopotamab deruxtecan?
 - a. HER2
 - b. Nectin-4
 - c. TIGIT
 - d. TROP2
 - e. CEACAM5
- 2. The Phase III REVEL trial evaluated the addition of ramucirumab to docetaxel versus docetaxel alone in which subgroup of patients with metastatic NSCLC?
 - a. Patients with nonsquamous histology
 - b. Patients with squamous histology
 - c. Both a and b
- 3. The ongoing Phase III Pragmatica-Lung (SWOG-S2302) study is evaluating which of the following agents in combination with pembrolizumab versus standard treatment for patients with Stage IV or recurrent NSCLC previously treated with immunotherapy?
 - a. Bevacizumab
 - b. Cabozantinib
 - c. Lenvatinib
 - d. Ramucirumab
 - e. Sitravatinib

- 4. Which of the following treatmentemergent adverse events was most prominent among patients treated with triplet therapy (Dato-DXd/pembrolizumab/ chemotherapy) on the TROPION-Lung02 trial?
 - a. Alopecia
 - b. Nausea
 - c. Neutropenia
 - d. Insomnia
- 5. Tumor treating fields promote antitumor activity by directly affecting which of the following cellular processes?
 - a. Cell division
 - b. Protein synthesis
 - c. Ion transport
 - d. Adhesion to matrix