POST-TEST

The Implications of Recent Datasets for the Current and Future Management of Breast Cancer — An ASCO 2025 Review

THE CORRECT ANSWER IS INDICATED WITH YELLOW HIGHLIGHTING.

- Which of the following measures is recommended to prevent or mitigate oral mucositis in patients receiving datopotamab deruxtecan (Dato-DXd)?
 - a. Systemic corticosteroids
 - b. Steroid mouthwash
 - c. Inhaled corticosteroids
 - d. No prophylactic measures are recommended when administering Dato-DXd
- 2. What is the maximum number of cycles of pembrolizumab a patient with metastatic triple-negative breast cancer may receive in the Saci-IO TNBC study?
 - a. 12 cycles
 - b. 24 cycles
 - c. 35 cycles
 - d. 48 cyclese. Indefinite
- 3. Which of the following adverse events
- is most commonly observed in patients receiving sacituzumab govitecan for previously treated advanced breast cancer?
 - a. Oral mucositis
 - b. Dry eye
 - c. Headaches
 - d. Diarrhea

- 4. Which of the following rare but serious adverse events has been observed in patients receiving Dato-DXd for previously treated advanced breast cancer?
 - a. Ventricular arrythmia
 - b. Deep vein thrombosis
 - c. Interstitial lung disease
 - d. Ischemic stroke
- 5. The DESTINY-Breast09 study evaluated trastuzumab deruxtecan for breast cancer in which setting?
 - With pembrolizumab for relapsed/ refractory HER2-low, PD-L1-positive advanced disease
 - b. With pertuzumab as first-line therapy for HER2-positive advanced disease
 - c. As adjuvant therapy for localized HER2-positive disease after complete resection