POST-TEST

Cases from the Community: Investigators Discuss Available Research Guiding the Care of Patients with Ovarian and Endometrial Cancer

THE CORRECT ANSWER IS INDICATED WITH YELLOW HIGHLIGHTING.

- 1. Raludotatug deruxtecan (R-DXd) is an antibody-drug conjugate directed against...
 - a. FRα
 - b. CDH6
 - c. HER2
 - d. XP01
- 2. The Phase III SIENDO trial revealed a statistically significant reduction in the risk of progression or death for patients with endometrial cancer (EC) receiving selinexor in which of the following settings?
 - a. As monotherapy after disease progression on first-line chemotherapy
 - b. As maintenance after induction chemotherapy
 - c. In combination with pembrolizumab for recurrent disease
 - d. As monotherapy after 3 or more prior lines of treatment
- 3. What was the approximate objective response rate with trastuzumab deruxtecan for patients with HER2 IHC 3+ advanced EC in the Phase II DESTINY-PanTumorO2 study?
 - a. 25%
 - b. 40%
 - c. 65%
 - d. 85%

- 4. Which of the following adverse events is most commonly associated with R-DXd?
 - a. Headache
 - b. Pneumonitis
 - c. Rash
 - d. Nausea
- 5. Which of the following descriptions best reflects results of the Phase III ROSELLA trial of relacorilant with *nab* paclitaxel for patients with platinum-resistant ovarian cancer?
 - a. No significant improvement in progression-free survival (PFS) or overall survival (OS)
 - b. Significant improvement in PFS only
 - c. Significant improvements in both PFS and OS