

THE CORRECT ANSWER IS INDICATED WITH YELLOW HIGHLIGHTING.

1. What was the approximate complete response rate in the ASCERTAIN-V trial of the all-oral regimen of decitabine/cedazuridine with venetoclax for patients with newly diagnosed acute myeloid leukemia (AML) not eligible for intensive induction chemotherapy?
 - a. 17%
 - b. 47%**
 - c. 87%
2. The ongoing Phase II QuANTUM-Wild study is evaluating quizartinib with intensive chemotherapy for which patients with newly diagnosed AML?
 - a. Those with FLT3-TKD-negative AML
 - b. Those with FLT3-ITD-negative AML**
 - c. Those with FLT3-TKD- and FLT3-ITD-negative AML
3. Which of the following conditions is a treatment-related adverse event of special interest associated with menin inhibitors?
 - a. Ocular toxicity
 - b. Differentiation syndrome**
 - c. Interstitial lung disease
4. Olutasidenib has demonstrated activity in which category of AML?
 - a. FLT3-ITD mutant
 - b. FLT3-TKD mutant
 - c. IDH1-mutant**
5. The BEAT AML trial of triplet therapy with azacitidine/venetoclax/revumenib for newly diagnosed KMT2A-rearranged or NPM1-mutant AML demonstrated which finding?
 - a. The overall response rate (ORR) in the overall patient population was approximately 88%**
 - b. Patients with KMT2A-rearranged AML did not respond to the triplet regimen
 - c. The ORR was higher for patients with NPM1-mutant AML than for those with KMT2A-rearranged AML